

2013/26



Universal gradient methods  
for convex optimization problems

Yu. Nesterov



**CORE**

DISCUSSION PAPER

Center for Operations Research  
and Econometrics

Voie du Roman Pays, 34  
B-1348 Louvain-la-Neuve  
Belgium

<http://www.uclouvain.be/core>

CORE DISCUSSION PAPER  
2013/26

**Universal gradient methods  
for convex optimization problems**

Yu. NESTEROV<sup>1</sup>

April 2013

**Abstract**

In this paper, we present new methods for black-box convex minimization. They do not need to know in advance the actual level of smoothness of the objective function. The only essential input parameter is the required accuracy of the solution. At the same time, for each particular problem class they automatically ensure the best possible rate of convergence. We confirm our theoretical results by encouraging numerical experiments, which demonstrate that the fast rate of convergence, typical for the smooth optimization problems, sometimes can be achieved even on nonsmooth problem instances.

**Keywords:** convex optimization, black-box methods, complexity bounds, optimal methods, weakly smooth functions

---

<sup>1</sup> Université catholique de Louvain, CORE, B-1348 Louvain-la-Neuve, Belgium. E-mail: yurii.nesterov@uclouvain.be. This author is also member of ECORE, the association between CORE and ECARES.

The research results presented in this paper have been supported by a grant "Action de recherche concertée ARC 04/09-315" from the "Direction de la recherche scientifique – Communauté française de Belgique".

This paper presents research results of the Belgian Program on Interuniversity Poles of Attraction initiated by the Belgian State, Prime Minister's Office, Science Policy Programming. The scientific responsibility is assumed by the author.

# 1 Introduction

**Motivation.** In Convex Optimization, the majority of numerical schemes are developed for particular problem classes. In the Black-Box framework, two main classes of convex problems, the smooth problems, and nonsmooth ones are treated by completely different techniques.

This separation looks very natural. Indeed, differentiable function allow constructing *monotone* minimization sequences, for which the convergence results can be easily obtained. Smooth function can be locally approximated by first- and second-order models, which are very helpful in developing efficient minimization schemes.

The class of nonsmooth convex functions looks much more difficult. For them, there is no hope to get a good local approximation model. It is very difficult to construct relaxation sequences. Moreover, even if a descent direction is found, there is no guarantee that we can advance along it by a sufficiently long step. Therefore, *all* methods for nonsmooth convex optimization rely only on separation properties. Cutting planes provide us with information about the half-spaces containing the optimal solution. Using this very restricted knowledge, it is still possible to develop some optimization methods. But their computational abilities are incomparably weaker than the abilities of smooth minimization schemes.

Above observations are confirmed by theoretical results. It is well known that for the class of smooth problems  $C^{1,1}(R^n)$ , composed by functions with Lipschitz-continuous gradients, the optimal iteration complexity bound for finding  $\epsilon$ -solution of corresponding optimization problem by a first-order method is of the order  $O(\frac{1}{\epsilon^{1/2}})$ . For nonsmooth problems from the class  $C^{1,0}(R^n)$ , where we can rely only on Lipschitz continuity of function values, such a bound is established on the level of  $O(\frac{1}{\epsilon^2})$  (see, e.g. [8]).

Such a big difference in the complexity bounds stimulated an interest to the intermediate classes of convex problems. One of the possibilities consists in considering functions from the class  $C^{1,\nu}(R^n)$ ,  $\nu \in [0, 1]$ , which have Hölder continuous gradients:

$$\|\nabla f(x) - \nabla f(y)\|_* \leq L_\nu \|x - y\|^\nu, \quad x, y \in R^n. \quad (1.1)$$

General Complexity Theory [7] established for this class the following lower iteration complexity bound:

$$O\left(\left(\frac{L_\nu R^{1+\nu}}{\epsilon}\right)^{\frac{2}{1+3\nu}}\right), \quad (1.2)$$

where  $R$  is the distance from a starting point to the solution. The first optimal methods for such problems were developed in [6]<sup>1</sup>. The main advantage of these schemes is an automatic adjustment to the proper level of smoothness parameter  $\nu$ . However, these methods need to know another characteristics of the problem (estimate of Lipschitz constant  $L_\nu$ , estimate of the distance to optimum), which are not readily available. Moreover, it was necessary to decide in advance on the total number of steps of the method. This requirement is not very practical. Indeed, in order to make such a decision, we need to know the rate of convergence of the method. However, this is possible only if we know the Hölder parameter. This hidden contradiction probably explains why these theoretically attractive procedures were never seriously tested in computational practice.

---

<sup>1</sup>English translation of this paper was included in Section 2.3 in [3]

In the last decade, we can see a restoration of interest to the gradient methods. New problems setting in image processing, data mining, and statistics require computationally cheap minimization procedures, which can quickly deliver an approximate solutions with a moderate accuracy. This demand was satisfied by new families of problem-oriented methods (e.g. [9], [10], [1]), which increase the rate of convergence of the gradient schemes much above the limits of Black-Box Complexity Theory [7]. This can be done, of-course, only by an appropriate use of problem structure, violating one of the main assumptions of the Black-Box concept.

However, it appears that the Black-Box methods did not reach yet the limits of their performance. The old idea of automatic adjustment to Hölder parameter was revived in [4], where a new version of Level Method [5] was adapted to smooth problems, ensuring the best possible complexity bounds for all values of the smoothness parameter. The only drawback of this approach is related to a high iteration cost of the Level Method.

Minimization of functions with Hölder-continuous gradient was discussed in [2] in the framework of *inexact oracle*. It was shown that the answer  $(f(x), \nabla f(x))$  of an *exact* oracle for a convex function satisfying Hölder condition (1.1) can be treated as “inexact” information for some function from  $C^{1,1}(R^n)$ :

$$0 \leq f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \tilde{\delta} + \frac{1}{2} \tilde{L} \|y - x\|^2, \quad x, y \in R^n, \quad (1.3)$$

where  $\tilde{L}$  and  $\tilde{\delta}$  are some “inexactness” parameters. It was shown that these parameters can be chosen as appropriate functions of  $\nu$ . Therefore, functions from  $C^{1,\nu}(R^n)$  can be minimized by an “inexact” version of Fast Gradient Methods for  $C^{1,1}(R^n)$ . The resulting complexity bounds appear to be optimal (1.2). However, in order to apply this approach, we need to employ a lot of additional information (the value of parameter  $\nu$ , constant  $L_\nu$ , distance estimate  $R$ , and the total number of steps of the method).

In this paper, we construct new universal methods for minimizing functions with Hölder-continuous gradient. They do not need à priori knowledge of the parameter  $\nu$ , and they have a cheap cost of one iteration.

In order to solve the problem

$$\min_{x \in Q} f(x) \quad (1.4)$$

by *universal* methods, we suggest to use a *Damped Relaxation Condition* (DR)

$$f(x_+) \leq \delta + \min_{y \in Q} \left[ f(\bar{x}) + \langle \nabla f(\bar{x}), y - \bar{x} \rangle + \frac{1}{2} \hat{L} \|y - \bar{x}\|^2 \right], \quad (1.5)$$

where the tolerance parameter  $\delta > 0$  depends only on the required *accuracy*  $\epsilon > 0$  of the final approximate solution. Similar conditions were used in [6] and [2] with  $\delta$  being a function of smoothness parameter  $\nu$  and total number of iterations. We show that all necessary information on  $\nu$  and  $L_\nu$  can be accumulated in the constant  $\hat{L}$ , which can be easily adapted by an appropriate “line-search” strategy. For different methods, the dependence of  $\delta$  in  $\epsilon$  must be different. For the simplest Primal and Dual Gradient Methods, it is enough to take  $\delta = \frac{\epsilon}{2}$ . For the Fast Gradient Method [9], we use condition (1.5) with much smaller value of  $\delta$ , allowing to maintain a damped version of the estimating sequence condition

$$A_k(f(x_k) - \frac{\epsilon}{2}) \leq \min_{x \in Q} \phi_k(x), \quad (1.6)$$

(see Section 4 for details).

All our methods are developed for composite minimization problems [10], which space of variables is endowed with arbitrary norm. Hence, we apply machinery of Bregman distances. Our methods adjust automatically to the actual level of smoothness of the smooth part of the objective function. The only essential input parameter for these schemes is the required accuracy  $\epsilon > 0$ .

**Contents.** The paper is organized as follows. In Section 2 we introduce the problem formulation and discuss the main properties of Bregman mapping as applied to functions with Hölder continuous gradients. After that, we prove a convergence result for Universal Primal Gradient Method and derive its complexity bound. We show that this method needs in average at most two calls of oracle per iteration. Moreover, this method can be equipped with a reliable stopping criterion.

In Section 3, we prove similar results for Universal Dual Gradient Method. This method needs in average four calls of oracle per iteration. Both these methods are based on DR-condition (1.5).

In Section 4, in order to derive Universal Fast Gradient Method, we introduce condition (1.6). We show that this scheme is uniformly optimal for minimizing composite function, which has Hölder-continuous gradients of its smooth part. This scheme has a reliable stopping criterion. It needs in average four calls of oracle per iteration.

In Section 5, we present preliminary computational results. We consider three families of random test problems. All of them are nonsmooth problems with Lipschitz-continuous objective function. It is shown that quite often the Universal Fast Gradient Method is able to accelerate and demonstrate the rate of convergence typical for smooth minimization schemes. The choice of appropriate norms is always very important.

**Notation.** In what follows, we work in a finite-dimensional linear vector space  $E$ . Its dual space, the space of all linear function on  $E$ , is denoted by  $E^*$ . For  $x \in E$  and  $s \in E^*$ , we denote by  $\langle s, x \rangle$  the value of linear function  $s$  at  $x$ . For the (primal) space  $E$ , we introduce a norm  $\|\cdot\|$ . Then the dual norm is defined in the standard way:

$$\|s\| \stackrel{\text{def}}{=} \max_{x \in E} \{\langle s, x \rangle : \|x\| \leq 1\}.$$

Finally, for a convex function  $f : \text{dom } f \rightarrow R$  with  $\text{dom } f \subseteq E$  we denote by  $\nabla f(x) \in E^*$  one of its subgradients.

## 2 Universal Primal Gradient Method

Consider the following minimization problem:

$$\min_{x \in Q} \left[ \tilde{f}(x) \stackrel{\text{def}}{=} f(x) + \Psi(x) \right], \quad (2.1)$$

where  $Q$  is a simple closed convex set,  $\Psi$  is a simple closed convex function. Function  $f$  is assumed to be subdifferentiable on  $Q$ . In order to characterize variability of its (sub)gradients, we introduce the following values:

$$M_\nu \equiv M_\nu(f) = \sup_{\substack{x, y \in Q, \\ x \neq y}} \frac{\|\nabla f(x) - \nabla f(y)\|_*}{\|x - y\|^\nu}, \quad \nu \geq 0. \quad (2.2)$$

This condition can be rewritten in the form

$$\ln M_\nu = \sup_{\substack{x, y \in Q, \\ x \neq y}} [ \ln \|\nabla f(x) - \nabla f(y)\|_* - \nu \ln \|x - y\| ].$$

Thus,  $M_\nu$  is a log-convex function of  $\nu$ . For certain  $\nu \in [0, 1]$ , the constant  $M_\nu$  can be infinite. However, if  $M_0$  and  $M_1$  are finite, then

$$M_\nu \leq M_0^{1-\nu} M_1^\nu, \quad 0 \leq \nu \leq 1. \quad (2.3)$$

In any case, if  $M_\nu < \infty$ , then

$$\|\nabla f(x) - \nabla f(y)\|_* \leq M_\nu \|x - y\|^\nu, \quad x, y \in Q. \quad (2.4)$$

This inequality ensures that

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M_\nu}{1+\nu} \|x - y\|^{1+\nu}, \quad x, y \in Q. \quad (2.5)$$

Our main assumption is as follows.

**Assumption 1**  $\hat{M}(f) \stackrel{\text{def}}{=} \inf_{0 \leq \nu \leq 1} M_\nu(f) < +\infty$ .

For solving problem (2.1), we introduce a *prox-function*  $d(x)$ . This is a differentiable strongly convex function with convexity parameter equal to one:

$$d(y) \geq d(x) + \langle \nabla d(x), y - x \rangle + \frac{1}{2} \|x - y\|^2, \quad x, y \in \text{rint } Q. \quad (2.6)$$

We assume that  $d(x)$  attains its minimum on  $Q$  at some point  $x_0$ , and  $d(x_0) = 0$ . Thus,

$$d(x) \stackrel{(2.6)}{\geq} \frac{1}{2} \|x - x_0\|^2, \quad x \in Q. \quad (2.7)$$

This prox-function defines the *Bregman distance*  $\xi(x, y) \stackrel{\text{def}}{=} d(y) - d(x) - \langle \nabla d(x), y - x \rangle$ . Clearly,  $\xi(x, x) \equiv 0$ , and

$$\xi(x, y) \stackrel{(2.6)}{\geq} \frac{1}{2} \|x - y\|^2, \quad x, y \in Q. \quad (2.8)$$

Now for any  $x \in Q$  we can define the *Bregman mapping*

$$\mathcal{B}_M(x) = \arg \min_{y \in Q} \left\{ \psi_M(x, y) \stackrel{\text{def}}{=} f(x) + \langle \nabla f(x), y - x \rangle + M \xi(x, y) + \Psi(y) \right\}. \quad (2.9)$$

We assume that this point is easily computable either in a closed form, or by some cheap computational procedure. The first-order optimality condition for optimization problem in (2.9) is as follows:

$$\langle \nabla f(x) + M(\nabla d(\mathcal{B}_M(x)) - \nabla d(x)) + \nabla \Psi(\mathcal{B}_M(x)), y - \mathcal{B}_M(x) \rangle \geq 0, \quad y \in Q. \quad (2.10)$$

Denote  $\psi_M^*(x) = \psi_M(x, \mathcal{B}_M(x))$ .

**Lemma 1** *Let function  $f$  satisfy condition (2.4). Then for any  $\delta > 0$  and*

$$M \geq \left[ \frac{1}{\delta} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}} \quad (2.11)$$

we have

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2} M \|y - x\|^2 + \frac{\delta}{2}, \quad x, y \in Q. \quad (2.12)$$

Therefore,

$$\tilde{f}(\mathcal{B}_M(x)) \leq \psi_M^*(x) + \frac{\delta}{2}. \quad (2.13)$$

**Proof:**

It is well known that for all nonnegative  $\tau$  and  $s$  we have

$$\frac{1}{p} \tau^p + \frac{1}{q} s^q \geq \tau s,$$

where  $p, q \geq 1$  and  $\frac{1}{p} + \frac{1}{q} = 1$ . Therefore, taking  $p = \frac{2}{1+\nu}$ ,  $q = \frac{2}{1-\nu}$ , and  $\tau = t^{1+\nu}$ , we get

$$t^{1+\nu} \leq \frac{1+\nu}{2s} t^2 + \frac{1-\nu}{2} s^{\frac{1+\nu}{1-\nu}}, \quad s > 0, t \geq 0.$$

Denote  $\delta = \frac{1-\nu}{1+\nu} M_\nu s^{\frac{1+\nu}{1-\nu}}$ . Then  $s = \left[ \frac{1+\nu}{1-\nu} \cdot \frac{\delta}{M_\nu} \right]^{\frac{1-\nu}{1+\nu}}$ . Therefore,

$$\frac{M_\nu}{1+\nu} t^{1+\nu} \leq \frac{1}{2s} M_\nu t^2 + \frac{\delta}{2} = \frac{1}{2} \left[ \frac{1+\nu}{1-\nu} \cdot \frac{\delta}{M_\nu} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}} t^2 + \frac{\delta}{2} \stackrel{(2.11)}{\leq} \frac{1}{2} M t^2 + \frac{\delta}{2}. \quad (2.14)$$

This inequality, together with (2.5), justifies (2.12). Further, denoting  $x_+ = \mathcal{B}_M(x)$ , we obtain:

$$\begin{aligned} f(x_+) &\stackrel{(2.5)}{\leq} f(x) + \langle \nabla f(x), x_+ - x \rangle + \frac{M_\nu}{1+\nu} \|x_+ - x\|^{1+\nu} \\ &\stackrel{(2.14)}{\leq} f(x) + \langle \nabla f(x), x_+ - x \rangle + \frac{M}{2} \|x_+ - x\|^2 + \frac{\delta}{2} \\ &\stackrel{(2.8)}{\leq} f(x) + \langle \nabla f(x), x_+ - x \rangle + M \xi(x, x_+) + \frac{\delta}{2}. \end{aligned}$$

Therefore,  $\tilde{f}(x_+) = f(x_+) + \Psi(x_+) \leq \psi_M^*(x) + \frac{\delta}{2}$ .  $\square$

Note that the right-hand side of inequality (2.11) is continuous in  $\nu$ . As  $\nu \rightarrow 1$ , it becomes

$$M \geq M_1. \quad (2.15)$$

Let us look now at the simplest Universal Primal Gradient Method equipped with a backtracking line search procedure with restore. Denote by  $x^*$  the optimal solution

to (2.1).

**Universal Primal Gradient Method (PGM)**

**Initialization.** Choose  $L_0 > 0$  and accuracy  $\epsilon > 0$ .

**For  $k \geq 0$  do:**

1. Find the smallest  $i_k \geq 0$  such that

$$\tilde{f}\left(\mathcal{B}_{2^{i_k}L_k}(x_k)\right) \leq \psi_{2^{i_k}L_k}^*(x_k) + \frac{1}{2}\epsilon.$$

2. Set  $x_{k+1} = \mathcal{B}_{2^{i_k}L_k}(x_k)$ , and  $L_{k+1} = 2^{i_k-1}L_k$ .

(2.16)

Denote  $\gamma(M, \epsilon) \stackrel{\text{def}}{=} \left[\frac{1}{\epsilon}\right]^{\frac{1-\nu}{1+\nu}} M^{\frac{2}{1+\nu}}$ ,  $S_k = \sum_{i=1}^{k+1} \frac{1}{L_k}$ , and  $\tilde{f}_k^* = \frac{1}{S_k} \sum_{i=0}^k \frac{1}{L_{i+1}} \tilde{f}(x_i)$ .

**Theorem 1** *Let  $f$  satisfies condition (2.4). Assume that  $L_0 \leq \gamma(M_\nu, \epsilon)$ . Then for all  $k \geq 0$  we have  $L_{k+1} \leq \gamma(M_\nu, \epsilon)$ . Moreover, for all  $y \in Q$*

$$\tilde{f}_k^* \leq \frac{1}{S_k} \sum_{i=0}^k \frac{1}{L_{i+1}} [f(x_i) + \langle \nabla f(x_i), y - x_i \rangle] + \Psi(y) + \frac{\epsilon}{2} + \frac{2}{S_k} \xi(x_0, y). \quad (2.17)$$

Therefore,  $\tilde{f}_k^* - \tilde{f}(x^*) \leq \frac{\epsilon}{2} + \frac{2\gamma(M_\nu, \epsilon)}{k+1} \xi(x_0, x^*)$ .

**Proof:**

In view of Lemma 1, the line-search procedure of Step 1 in method (2.16) is well defined, and

$$2L_{k+1} = 2^{i_k}L_k \leq 2\gamma(M_\nu, \epsilon). \quad (2.18)$$

Let us fix an arbitrary point  $y \in Q$ . Denote  $r_k(y) \stackrel{\text{def}}{=} \xi(x_k, y)$ . Then

$$\begin{aligned} r_{k+1}(y) &= d(y) - d(x_{k+1}) - \langle \nabla d(x_{k+1}), y - x_{k+1} \rangle \\ &\stackrel{(2.10)}{\leq} d(y) - d(x_{k+1}) - \langle \nabla d(x_k), y - x_{k+1} \rangle + \frac{1}{2L_{k+1}} \langle \nabla f(x_k) + \nabla \Psi(x_{k+1}), y - x_{k+1} \rangle. \end{aligned}$$

Note that

$$\begin{aligned} &d(y) - d(x_{k+1}) - \langle \nabla d(x_k), y - x_{k+1} \rangle \\ &\stackrel{(2.6)}{\leq} d(y) - d(x_k) - \langle \nabla d(x_k), x_{k+1} - x_k \rangle - \frac{1}{2} \|x_{k+1} - x_k\|^2 - \langle \nabla d(x_k), y - x_{k+1} \rangle \\ &= r_k(y) - \frac{1}{2} \|x_{k+1} - x_k\|^2. \end{aligned}$$



Thus,

$$\begin{aligned}
r_{k+1}(y) - r_k(y) &\leq \frac{1}{2L_{k+1}} \langle \nabla f(x_k) + \nabla \Psi(x_{k+1}), y - x_{k+1} \rangle - \frac{1}{2} \|x_{k+1} - x_k\|^2 \\
&= \frac{1}{2L_{k+1}} \langle \nabla \Psi(x_{k+1}), y - x_{k+1} \rangle - \frac{1}{2L_{k+1}} (\langle \nabla f(x_k), x_{k+1} - x_k \rangle + L_{k+1} \|x_{k+1} - x_k\|^2) \\
&\quad + \frac{1}{2L_{k+1}} \langle \nabla f(x_k), y - x_k \rangle \\
&\leq \frac{1}{2L_{k+1}} \left( \Psi(y) - \Psi(x_{k+1}) + f(x_k) - f(x_{k+1}) + \frac{1}{2} \epsilon + \langle \nabla f(x_k), y - x_k \rangle \right).
\end{aligned}$$

Thus, we obtain the following inequality:

$$\frac{1}{2L_{k+1}} \tilde{f}(x_{k+1}) + r_{k+1}(y) \leq \frac{1}{2L_{k+1}} \left( f(x_k) + \langle \nabla f(x_k), y - x_k \rangle + \Psi(y) + \frac{\epsilon}{2} \right) + r_k(y).$$

Summing up these inequalities, we obtain

$$\tilde{f}_k^* \leq \frac{1}{S_k} \sum_{i=0}^k \frac{1}{L_{i+1}} [f(x_i) + \langle \nabla f(x_i), y - x_i \rangle + \Psi(y) + \frac{\epsilon}{2} + \frac{2}{S_k} r_0(y)].$$

It remains to use inequality (2.18). □

It is important that method (2.16) does not include  $\nu$  as a parameter. Therefore, in view of Theorem (1), in order to get  $\epsilon$ -solution of problem (2.1) we need

$$4\xi(x_0, x^*) \inf_{0 \leq \nu \leq 1} \left( \frac{M_\nu}{\epsilon} \right)^{\frac{2}{1+\nu}} \quad (2.19)$$

iterations at most. In this estimate, among all classes of functions with Hölder continuous gradient, we choose the class which better fits our particular objective function. Note that the expression (2.19) is log-quasiconvex in  $\nu$ . Hence, if  $M_0$  and  $M_1$  are finite, there are good chances that the optimal  $\nu$  belongs to the interval  $(0, 1)$ .

Inequality (2.17) gives us a reliable stopping criterion for method (2.16). Indeed, assume we have a bound for the size of optimal solution:

$$\xi(x_0, x^*) \leq D. \quad (2.20)$$

Denote  $\ell_k^p(y) \stackrel{\text{def}}{=} \frac{1}{S_k} \sum_{i=0}^k \frac{1}{L_{i+1}} [f(x_i) + \langle \nabla f(x_i), y - x_i \rangle]$ , and define

$$\hat{f}_k = \min_{y \in Q} \{ \ell_k^p(y) + \Psi(y) : \xi(x_0, y) \leq D \}.$$

Then

$$\tilde{f}_k^* - \tilde{f}(x^*) \leq \tilde{f}_k^* - \hat{f}_k \leq \frac{2\gamma(M_\nu, \epsilon)}{k+1} D. \quad (2.21)$$

Note that  $\hat{f}_k$  can be computed. Thus, inequality (2.21) provides us with an implementable stopping criterion  $\tilde{f}_k^* - \hat{f}_k \leq \epsilon$ .

Finally, let us estimate  $N(k)$ , the total number of computations of the function values in method (2.16) after  $k$  iterations. Note that

$$L_{k+1} = \frac{1}{2} 2^{i_k} L_k.$$

Therefore,  $i_k - 1 = \log_2 \frac{L_{k+1}}{L_k}$ . Hence, for any  $\nu \in [0, 1]$ , we have

$$\begin{aligned} N(k) &= \sum_{j=0}^k (i_j + 1) = 2(k+1) + \log_2 L_{k+1} - \log_2 L_0 \\ &\stackrel{(2.18)}{\leq} 2(k+1) + \frac{1-\nu}{1+\nu} \log_2 \frac{1}{\epsilon} + \frac{2}{1+\nu} \log_2 M_\nu - \log_2 L_0. \end{aligned}$$

Finally, we come to the following upper bound:

$$N(k) \leq 2(k+1) - \log_2 L_0 + \inf_{0 \leq \nu \leq 1} \left[ \frac{1-\nu}{1+\nu} \log_2 \frac{1}{\epsilon} + \frac{2}{1+\nu} \log_2 M_\nu \right]. \quad (2.22)$$

Thus in average, up to negligible logarithmic terms, method (2.16) requires two computations of function values per iteration.

The complexity estimates in (2.19) are optimal only for  $\nu = 0$ . In Section 4 we show that much better (and optimal) bounds can be achieved by a fast gradient scheme.

### 3 Universal Dual Gradient Method

Dual gradient method is based on updating a simple model for objective function of problem (2.1). Its justification is based on the following simple result.

**Lemma 2** *Let  $\phi : Q \rightarrow R \cup \{+\infty\}$  be a convex function such that for some  $M \geq 0$  the difference  $\phi(x) - Md(x)$  is subdifferentiable on  $Q$ . Denote  $\bar{x} = \arg \min_{x \in Q} \phi(x)$ . Then*

$$\phi(y) \geq \phi(\bar{x}) + M\xi(\bar{x}, y), \quad y \in Q. \quad (3.1)$$

**Proof:**

Denote  $F(y) = \phi(y) - Md(y)$ . Then  $\langle \nabla F(\bar{x}), y - \bar{x} \rangle \geq 0$  for all  $y \in Q$ . Therefore,

$$\begin{aligned} \phi(y) &= F(y) + Md(y) \geq F(\bar{x}) + \langle \nabla F(\bar{x}), y - \bar{x} \rangle + Md(y) \\ &\geq F(\bar{x}) - M\langle \nabla d(\bar{x}), y - \bar{x} \rangle + Md(y) = \phi(\bar{x}) + M\xi(\bar{x}, y). \quad \square \end{aligned}$$

**Universal Dual Gradient Method (DGM)**

**Initialization.** Choose  $L_0 > 0$ . Define  $\phi_0(x) = \xi(x_0, x)$ .

**For  $k \geq 0$  do:**

1. Find the smallest  $i_k \geq 0$  such that for point

$$x_{k,i_k} = \arg \min_{x \in Q} \left\{ \phi_k(x) + \frac{1}{2^{i_k} L_k} [f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \Psi(x)] \right\} \quad (3.2)$$

we have  $\tilde{f}(\mathcal{B}_{2^{i_k} L_k}(x_{k,i_k})) \leq \psi_{2^{i_k} L_k}^*(x_{k,i_k}) + \frac{\epsilon}{2}$ .

2. Set  $x_{k+1} = x_{k,i_k}$ ,  $L_{k+1} = 2^{i_k-1} L_k$ , and

$$\phi_{k+1}(x) = \phi_k(x) + \frac{1}{2L_{k+1}} [f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \Psi(x)].$$

Assume that  $L_0 \leq \gamma(M_\nu, \epsilon)$ , and  $M_\nu < \infty$ . Note that the termination criterion of Step 1 in method (3.2) is satisfied if  $2^{i_k} L_k \geq \gamma(M_\nu, \epsilon)$ . Therefore, same as in the proof of Theorem 1, we have

$$L_k \leq \gamma(M_\nu, \epsilon), \quad k \geq 1. \quad (3.3)$$

Denote  $y_k = \mathcal{B}_{2^{i_k} L_k}(x_{k,i_k})$ ,  $S_k = \sum_{i=0}^k \frac{1}{L_{i+1}}$ , and  $\phi_k^* = \arg \min_{y \in Q} \phi_k(y)$ . Let us prove by induction that the relation

$$\sum_{i=0}^k \frac{1}{2L_{i+1}} \tilde{f}(y_i) \leq \phi_{k+1}^* + S_k \cdot \frac{\epsilon}{4} \quad (3.4)$$

is valid for all  $k \geq 0$ . Indeed, for  $k = 0$  we have

$$\begin{aligned} \frac{1}{2L_1} \tilde{f}(y_0) - S_0 \cdot \frac{\epsilon}{4} &= \frac{1}{2L_1} \left[ \tilde{f}(y_0) - \frac{\epsilon}{2} \right] \leq \frac{1}{2^{i_0} L_0} \psi_{2^{i_0} L_0}^*(y_0) \\ &= \frac{1}{2^{i_0} L_0} [f(x_0) + \langle \nabla f(x_0), y_0 - x_0 \rangle + \Psi(y_0)] + \xi(x_0, y_0) \\ &= \phi_1(y_0) = \min_{x \in Q} \phi_1(x). \end{aligned}$$

In view of Lemma 2, for any  $k \geq 0$  we have

$$\phi_k(x) \geq \phi_k(x_k) + \xi(x_k, x), \quad x \in Q.$$

Assume that (3.4) is true for some  $k \geq 0$ . Then,

$$\begin{aligned}
& \min_{x \in Q} \phi_{k+2}(x) \\
& \geq \min_{x \in Q} \left\{ \phi_{k+1}(x) + \frac{1}{2L_{k+2}} [f(x_{k+1}) + \langle \nabla f(x_{k+1}), x - x_{k+1} \rangle + \Psi(x)] \right\} \\
& \geq \min_{x \in Q} \left\{ \phi_{k+1}(x_{k+1}) + \xi(x_{k+1}, x) + \frac{1}{2L_{k+2}} [f(x_{k+1}) + \langle \nabla f(x_{k+1}), x - x_{k+1} \rangle + \Psi(x)] \right\} \\
& \geq \phi_{k+1}(x_{k+1}) + \frac{1}{2L_{k+2}} \left[ \tilde{f}(y_{k+1}) - \frac{\epsilon}{2} \right] \stackrel{(3.4)}{\geq} -S_{k+1} \cdot \frac{\epsilon}{4} + \sum_{i=0}^{k+1} \frac{1}{2L_{i+1}} \tilde{f}(y_i).
\end{aligned}$$

Thus, we have proved that (3.4) is valid for all  $k \geq 0$ . Now we can prove the main convergence result for Universal Dual Gradient Method. Denote  $\tilde{f}_k^* = \frac{1}{S_k} \sum_{i=0}^k \frac{1}{L_{i+1}} \tilde{f}(y_i)$ .

**Theorem 2** *Let  $f$  satisfies condition (2.4) with  $M_\nu < \infty$ , and  $L_0 \leq \gamma(M_\nu, \epsilon)$ . Then all  $L_k$  generated by method (3.2) satisfy condition (3.3). Moreover, for all  $k \geq 0$  we have*

$$\tilde{f}_k^* - \tilde{f}(x^*) \leq \frac{\epsilon}{2} + \frac{2\gamma(M_\nu, \epsilon)}{k+1} \xi(x_0, x^*). \quad (3.5)$$

**Proof:**

Indeed, in view of inequality (3.4), we have

$$\frac{1}{2} S_k \tilde{f}_k^* = \sum_{i=0}^k \frac{1}{2L_{i+1}} \tilde{f}(y_i) \leq \min_{x \in Q} \phi_{k+1}(x) + S_k \cdot \frac{\epsilon}{4} \leq \frac{1}{2} S_k \tilde{f}(x^*) + \xi(x_0, x^*) + S_k \cdot \frac{\epsilon}{4}.$$

It remains to use inequality (3.3).  $\square$

Note that the worst-case complexity bound for the number of iterations of method (3.2) coincides with the bound (2.19). However, the number of function evaluations at each iteration of (3.2) is twice more than (2.22).

Same as method (2.16), Universal Dual Gradient Method can be equipped with a stopping criterion. Denote  $\ell_k^d(y) = \sum_{i=0}^k \frac{1}{L_{i+1}} [f(x_i) + \langle \nabla f(x_i), y - x_i \rangle]$ . Assume that  $\xi(x_0, x^*) \leq D$  and the constant  $D$  is known. Denote

$$\hat{f}_k = \min_{y \in Q} \left\{ \frac{1}{S_k} \ell_k^d(y) + \Psi(y) : \xi(x_0, y) \leq D \right\}.$$

Note that

$$\begin{aligned}
\hat{f}_k &= \min_{x \in Q} \max_{\beta \geq 0} \left\{ \frac{1}{S_k} \ell_k^d(y) + \Psi(y) + \beta (\xi(x_0, y) - D) \right\} \\
&= \max_{\beta \geq 0} \min_{x \in Q} \left\{ \frac{1}{S_k} \ell_k^d(y) + \Psi(y) + \beta (\xi(x_0, y) - D) \right\} \\
&\stackrel{\beta=2/S_k}{\geq} \frac{2}{S_k} \phi_{k+1}^* - \frac{2}{S_k} D.
\end{aligned}$$

Since  $\tilde{f}_k^* \stackrel{(3.4)}{\leq} \frac{2}{S_k} \phi_{k+1}^* + \frac{\epsilon}{2}$ , we conclude that the stopping criterion  $\tilde{f}_k^* - \hat{f}_k \leq \epsilon$  ensures  $\tilde{f}_k^* - \tilde{f}(x^*) \leq \epsilon$  as far as  $S_k \geq \frac{4}{\epsilon} D$ .

## 4 Universal Fast Gradient Method

Finally, let us apply to problem (2.1) the following method.

<b>Universal Fast Gradient Method (FGM)</b>	
<p><b>Initialization.</b> Choose <math>L_0 &gt; 0</math>. Define <math>\phi_0(x) = \xi(x_0, x)</math>, <math>y_0 = x_0</math>, <math>A_0 = 0</math>.</p> <p><b>For <math>k \geq 0</math> do:</b></p> <ol style="list-style-type: none"> <li>1. Find <math>v_k = \arg \min_{x \in Q} \phi_k(x)</math>.</li> <li>2. Find the smallest <math>i_k \geq 0</math> such that coefficient <math>a_{k+1, i_k} &gt; 0</math>, computed from equation <math>a_{k+1, i_k}^2 = \frac{1}{2^{i_k} L_k} (A_k + a_{k+1, i_k})</math> and used in the definitions</li> </ol> $A_{k+1, i_k} = A_k + a_{k+1, i_k}, \quad \tau_{k, i_k} = \frac{a_{k+1, i_k}}{A_{k+1, i_k}}, \quad x_{k+1, i_k} = \tau_{k, i_k} v_k + (1 - \tau_{k, i_k}) y_k, \quad (4.1)$ $\hat{x}_{k+1, i_k} = \arg \min_{y \in Q} \{ \xi(v_k, y) + a_{k+1, i_k} [\langle \nabla f(x_{k+1, i_k}), y \rangle + \Psi(y)] \},$ $y_{k+1, i_k} = \tau_{k, i_k} \hat{x}_{k+1, i_k} + (1 - \tau_{k, i_k}) y_k, \text{ ensures the following relation:}$ $f(y_{k+1, i_k}) \leq f(x_{k+1, i_k}) + \langle \nabla f(x_{k+1, i_k}), y_{k+1, i_k} - x_{k+1, i_k} \rangle + 2^{i_k - 1} L_k \ y_{k+1, i_k} - x_{k+1, i_k}\ ^2 + \frac{\epsilon}{2} \tau_{k, i_k}.$ <ol style="list-style-type: none"> <li>3. Set <math>x_{k+1} = x_{k+1, i_k}</math>, <math>y_{k+1} = y_{k+1, i_k}</math>, <math>a_{k+1} = a_{k+1, i_k}</math>, <math>\tau_k = \tau_{k, i_k}</math>.</li> </ol> <p>Define <math>A_{k+1} = A_k + a_{k+1}</math>, <math>L_{k+1} = 2^{i_k - 1} L_k</math>, and</p> $\phi_{k+1}(x) = \phi_k(x) + a_{k+1} [f(x_{k+1}) + \langle \nabla f(x_{k+1}), x - x_{k+1} \rangle + \Psi(x)].$	

**Theorem 3** *Let  $f$  satisfies condition (2.4) with certain  $M_\nu < +\infty$ . Then all iterations of method (4.1) are well defined. Moreover, for all  $k \geq 0$  we have*

$$A_k \left( \tilde{f}(y_k) - \frac{\epsilon}{2} \right) \leq \phi_k^* \stackrel{\text{def}}{=} \min_{x \in Q} \phi_k(x), \quad (4.2)$$

where  $A_k \geq \left[ \frac{1}{2^{2+4\nu} M_\nu^2} \epsilon^{1-\nu} k^{1+3\nu} \right]^{\frac{1}{1+\nu}}$ . Consequently, for all  $k \geq 1$  we have

$$\tilde{f}(y_k) - \tilde{f}(x^*) \leq \left[ \frac{2^{2+4\nu} M_\nu^2}{\epsilon^{1-\nu} k^{1+3\nu}} \right]^{\frac{1}{1+\nu}} \xi(x_0, x^*) + \frac{\epsilon}{2}. \quad (4.3)$$

**Proof:**

Let us prove first, that the "line-search" process of Item 2 in (4.1) is finite. In view of inequality (2.12), we need to show that

$$2^{i_k} L_k \geq \left[ \frac{1}{\epsilon \tau_{k, i_k}} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}}$$

for  $i_k$  large enough. Indeed, in view of the characteristic equation for  $a_{k+1, i_k}$ , we have

$$2^{i_k} L_k \tau_{k, i_k}^{\frac{1-\nu}{1+\nu}} = \frac{A_{k+1, i_k}}{a_{k+1, i_k}^2} \cdot \left( \frac{a_{k+1, i_k}}{A_{k+1, i_k}} \right)^{\frac{1-\nu}{1+\nu}} = \left( \frac{1}{\tau_{k, i_k}} \right)^{\frac{2\nu}{1+\nu}} \cdot \frac{1}{a_{k+1, i_k}} \geq \frac{1}{a_{k+1, i_k}}.$$

It remains to note that  $a_{k+1, i_k} \rightarrow 0$  as  $i_k \rightarrow \infty$ .

Let us prove relation (4.2). For  $k = 0$  it is evident. Assume that it is valid for certain  $k \geq 0$ . Then for any  $y \in Q$  we have

$$\begin{aligned} \phi_k(y) &\stackrel{(3.1)}{\geq} \phi_k^* + \xi(v_k, y) \stackrel{(4.2)}{\geq} A_k \left( \tilde{f}(y_k) - \frac{\epsilon}{2} \right) + \xi(v_k, y_k) \\ &\geq A_k (f(x_{k+1}) + \langle \nabla f(x_{k+1}), y_k - x_{k+1} \rangle + \Psi(y_k) - \frac{\epsilon}{2}) + \xi(v_k, y). \end{aligned}$$

Therefore,

$$\begin{aligned} \phi_{k+1}(y) &\geq \xi(v_k, y) + A_k (f(x_{k+1}) + \langle \nabla f(x_{k+1}), y_k - x_{k+1} \rangle + \Psi(y_k) - \frac{\epsilon}{2}) \\ &\quad + a_{k+1} [f(x_{k+1}) + \langle \nabla f(x_{k+1}), y - x_{k+1} \rangle + \Psi(y)] \\ &= \xi(v_k, y) + A_k (f(x_{k+1}) + \Psi(y_k) - \frac{\epsilon}{2}) \\ &\quad + a_{k+1} [f(x_{k+1}) + \langle \nabla f(x_{k+1}), y - v_k \rangle + \Psi(y)]. \end{aligned}$$

In view of definition of point  $\hat{x}_{k+1}$ , we have

$$\begin{aligned} \phi_{k+1}^* &\geq \xi(v_k, \hat{x}_{k+1}) + A_k (f(x_{k+1}) + \Psi(y_k) - \frac{\epsilon}{2}) \\ &\quad + a_{k+1} [f(x_{k+1}) + \langle \nabla f(x_{k+1}), \hat{x}_{k+1} - v_k \rangle + \Psi(\hat{x}_{k+1})] \\ &\stackrel{(2.8)}{\geq} \frac{1}{2} \|\hat{x}_{k+1} - v_k\|^2 + A_{k+1} f(x_{k+1}) + A_{k+1} \Psi(y_{k+1}) - \frac{\epsilon}{2} A_k \\ &\quad + a_{k+1} \langle \nabla f(x_{k+1}), \hat{x}_{k+1} - v_k \rangle. \end{aligned}$$

Since  $\hat{x}_{k+1} - v_k = \frac{1}{\tau_k} (y_{k+1} - x_{k+1})$ , we obtain

$$\begin{aligned} \phi_{k+1}^* &\geq \frac{1}{2\tau_k^2} \|y_{k+1} - x_{k+1}\|^2 + A_{k+1} f(x_{k+1}) + A_{k+1} \Psi(y_{k+1}) - \frac{\epsilon}{2} A_k \\ &\quad + A_{k+1} \langle \nabla f(x_{k+1}), y_{k+1} - x_{k+1} \rangle \\ &= A_{k+1} (f(x_{k+1}) + \langle \nabla f(x_{k+1}), y_{k+1} - x_{k+1} \rangle) \\ &\quad + 2^{i_k-1} L_k \|y_{k+1} - x_{k+1}\|^2 + \Psi(y_{k+1}) - \frac{\epsilon}{2} A_k \\ &\geq A_{k+1} (f(y_{k+1}) - \frac{\epsilon}{2} \tau_k + \Psi(y_{k+1})) - \frac{\epsilon}{2} A_k = A_{k+1} \left( \tilde{f}(y_{k+1}) - \frac{\epsilon}{2} \right). \end{aligned}$$

Thus, inequality (4.2) is proved for all  $k \geq 0$ . Since  $\phi_k(y) \leq A_k \tilde{f}(y) + \xi(x_0, y)$  for all  $y \in Q$ , we obtain

$$\tilde{f}(y_k) - \tilde{f}(x^*) \stackrel{(4.2)}{\leq} \frac{\xi(x_0, x^*)}{A_k} + \frac{\epsilon}{2}, \quad k \geq 1.$$

It remains to estimate the growth of coefficients  $A_k$ .

In view of Lemma 1, the number of internal steps  $i_k$  in Item 2 of (4.1) satisfies inequality

$$2^{i_k} L_k \leq 2 \left[ \frac{1}{\epsilon \tau_k} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}}.$$

Therefore,  $\frac{a_{k+1}^2}{A_{k+1}} = \frac{1}{2^{i_k} L_k} \geq \frac{1}{2M_\nu^{\frac{2}{1+\nu}}} [\epsilon \tau_k]^{\frac{1-\nu}{1+\nu}}$ , which is  $a_{k+1}^2 \geq \frac{[\epsilon a_{k+1}]^{\frac{1-\nu}{1+\nu}}}{2M_\nu^{\frac{2}{1+\nu}}} A_{k+1}^{\frac{2\nu}{1+\nu}}$ . Thus, we come to the following estimate:

$$a_{k+1} \geq \frac{\epsilon^{\frac{1-\nu}{1+3\nu}} A_{k+1}^{\frac{2\nu}{1+3\nu}}}{2^{\frac{1+\nu}{1+3\nu}} M_\nu^{\frac{2}{1+3\nu}}}. \quad (4.4)$$

Denote  $B_k = A_k^\gamma$ , where  $\gamma = \frac{1+\nu}{1+3\nu} \geq \frac{1}{2}$ . Since  $A_{k+1} \geq A_k$ , we have

$$B_{k+1} - B_k \geq \frac{A_{k+1} - A_k}{A_{k+1}^{1-\gamma} + A_k^{1-\gamma}} \geq \frac{a_{k+1}}{2A_{k+1}^{1-\gamma}} \stackrel{(4.4)}{\geq} \frac{\epsilon^{\frac{1-\nu}{1+3\nu}}}{2^{\frac{2+4\nu}{1+3\nu}} M_\nu^{\frac{2}{1+3\nu}}}.$$

Thus, we have proved that  $A_k \geq \left[ \frac{k \cdot \epsilon^{\frac{1-\nu}{1+3\nu}}}{2^{\frac{2+4\nu}{1+3\nu}} M_\nu^{\frac{2}{1+3\nu}}} \right]^{\frac{1+3\nu}{1+\nu}} = \frac{k^{\frac{1+3\nu}{1+\nu}} \epsilon^{\frac{1-\nu}{1+\nu}}}{2^{\frac{2+4\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}}}$ .  $\square$

From the rate of convergence (4.3), we get the following upper bound for the number of iterations, which are necessary for getting  $\epsilon$ -solution of problem (2.1):

$$k \leq \inf_{0 \leq \nu \leq 1} \left[ \left( \frac{2^{\frac{3+5\nu}{2}} M_\nu}{\epsilon} \right)^{\frac{2}{1+3\nu}} \xi(x_0, x^*)^{\frac{1+\nu}{1+3\nu}} \right]. \quad (4.5)$$

As compared with (2.19), the dependence of this bound in smoothness parameters is now optimal.

Same as the gradient methods (2.16) and (3.2), Fast Gradient Method (4.1) can be equipped with an implementable stopping criterion. Assume that  $\xi(x_0, x^*) \leq D$ . Denote  $\ell_k^{pd}(y) = \sum_{i=1}^k a_i [f(x_i) + \nabla f(x_i), x - x_i]$ , and  $\hat{f}_k = \min_{y \in Q} \{ \frac{1}{A_k} \ell_k^{pd}(y) + \Psi(y) : \xi(x_0, y) \leq D \}$ .

Note that  $\tilde{f}(y_k) \stackrel{(4.2)}{\leq} \frac{\epsilon}{2} + \frac{1}{A_k} \phi_k^*$ . Using the reasoning presented in the end of Section 3, we obtain

$$\hat{f}_k = \max_{\beta \geq 0} \min_{y \in Q} \left\{ \frac{1}{A_k} \ell_k^{pd}(y) + \Psi(y) + \beta (\xi(x_0, y) - D) \right\} \stackrel{\beta=1/A_k}{\geq} \frac{1}{A_k} \phi_k^* - \frac{1}{A_k} D.$$

Thus, we can use stopping criterion

$$\tilde{f}(y_k) - \hat{f}_k \leq \frac{\epsilon}{2}, \quad (4.6)$$

which ensures  $\tilde{f}(y_k) - \tilde{f}(x^*) \leq \epsilon$  as far as

$$A_k \geq \frac{2}{\epsilon} D. \quad (4.7)$$

It remains to estimate from above the total number of calls of oracle of method (4.1), which is sufficient to get an  $\epsilon$ -solution of problem (2.1). Let us assume that this method is equipped with the stopping criterion (4.6). Then we can be sure that

$$A_k \leq \frac{2}{\epsilon} D, \quad k \geq 0. \quad (4.8)$$

Denote by  $N(k)$  the total number of calls of oracle after  $k$  iterations. At each iteration of this method we call the oracle  $2(i_k + 1)$  times (at point  $x_{k+1, i_k}$  and at the prediction point  $y_{k+1, i_k}$ ). Therefore, using the same reasoning as in the end of Section 2, we conclude that

$$N(k) = 4(k + 1) + 2 \log_2 L_{k+1} - 2 \log_2 L_0. \quad (4.9)$$

Note that

$$L_{k+1} = \frac{1}{2} 2^{i-k} L_k = \frac{A_{k+1}}{a_{k+1}^2} \stackrel{(2.11)}{\leq} \left[ \frac{1}{\epsilon 7^k} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}} = \left[ \frac{A_{k+1}}{\epsilon a_{k+1}} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}} \quad (4.10)$$

Therefore  $\left[ \frac{1}{a_{k+1}} \right]^{\frac{1+3\nu}{1+\nu}} \leq A_{k+1}^{\frac{-2\nu}{1+\nu}} \left[ \frac{1}{\epsilon} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}}$ , and we conclude that

$$\begin{aligned} L_{k+1} &\leq A_{k+1} \left[ A_{k+1}^{\frac{-2\nu}{1+\nu}} \left[ \frac{1}{\epsilon} \right]^{\frac{1-\nu}{1+\nu}} M_\nu^{\frac{2}{1+\nu}} \right]^{\frac{2(1+\nu)}{1+3\nu}} = A_{k+1}^{\frac{1-\nu}{1+3\nu}} \left[ \frac{1}{\epsilon} \right]^{\frac{2(1-\nu)}{1+3\nu}} M_\nu^{\frac{4}{1+3\nu}} \\ &\stackrel{(4.8)}{\leq} (2D)^{\frac{1-\nu}{1+3\nu}} \left[ \frac{1}{\epsilon} \right]^{\frac{3(1-\nu)}{1+3\nu}} M_\nu^{\frac{4}{1+3\nu}}. \end{aligned}$$

Substituting this estimate in the expression (4.9), we obtain that in average method (4.1) has at most four calls of oracle per iteration.

## 5 Numerical experiments

In our numerical experiments we tried to check the actual level of adaptivity of the above methods to the local topological structure of the objective function. For that, we have chosen three families of nonsmooth minimization problems.

**1. Matrix games.** In this problem, given by an  $n \times m$  payoff matrix  $A$ , we need to find a saddle point of the following problem:

$$\begin{aligned} \min_{x \in \Delta_n} \max_{y \in \Delta_m} \langle x, Ay \rangle &= \min_{x \in \Delta_n} \left\{ \psi_p(x) \stackrel{\text{def}}{=} \max_{1 \leq j \leq m} \langle x, Ae_j \rangle \right\} \\ &= \max_{y \in \Delta_m} \left\{ \psi_d(y) \stackrel{\text{def}}{=} \min_{1 \leq i \leq n} \langle e_i, Ay \rangle \right\}, \end{aligned} \quad (5.1)$$

where  $e_{(\cdot)}$  denote the basis vectors in the corresponding spaces, and  $\Delta_{(\cdot)}$  denotes the standard simplex. This problem can be posed as a minimization problem

$$\min_{x \in \Delta_n, y \in \Delta_m} \{ \psi_{pd}(x, y) = \psi_p(x) - \psi_d(y) \}. \quad (5.2)$$



The optimal value of this problem is zero. For our experiments, we generated matrix  $A$  randomly, with uniform distribution of its entries in the interval  $[-1, 1]$ .

For feasible set of this problem,  $\mathcal{F} = \{z = (x, y) : x \in \Delta_n, y \in \Delta_m\}$ , a natural prox-function is the *entropy*:

$$\eta(z) = \sum_{i=1}^n z^{(i)} \ln z^{(i)}.$$

This function is strongly convex with respect to  $\ell_1$ -norm with the convexity parameter one. Note that  $\ell_1$ -norm is very good for measuring simplexes. Consequently, we can measure the subgradients of the objective function in (5.2) in  $\ell_\infty$ -norm. In view of our strategy for generating the matrix  $A$ , we get Lipschitz-continuous function  $\psi_{pd}$  with the constant equal to one.

We will refer to the methods based on the entropy function as methods with the *Entropy Setup*. If a method is using the standard Euclidean norm, we say that it is based on the Euclidean setup.

In the table below, we give computational results for two universal methods, the Fast Gradient Method (4.1), and the Primal Gradient Method (2.16), both with Entropy Setup. In our problem instance,  $n = 896$  and  $m = 128$ .

$Eps$	$FGM_{Entropy}$			$PGM_{Entropy}$		
$2^{-5}$	516	$6.0 \cdot 10^{-2}$	$1.3 \cdot 10^2$	722	$8.2 \cdot 10^{-2}$	8.0
$2^{-6}$	1127	$2.9 \cdot 10^{-2}$	$2.6 \cdot 10^2$	2065	$5.2 \cdot 10^{-2}$	$1.6 \cdot 10^1$
$2^{-7}$	1937	$1.6 \cdot 10^{-2}$	$2.0 \cdot 10^2$	5675	$3.4 \cdot 10^{-2}$	$3.2 \cdot 10^1$
$2^{-8}$	4684	$7.9 \cdot 10^{-3}$	$2.0 \cdot 10^3$	15731	$2.3 \cdot 10^{-2}$	$6.4 \cdot 10^1$
$2^{-9}$	8129	$3.8 \cdot 10^{-3}$	$8.2 \cdot 10^3$	44829	$1.5 \cdot 10^{-2}$	$1.3 \cdot 10^2$
$2^{-10}$	17556	$2.1 \cdot 10^{-3}$	$4.1 \cdot 10^3$	122959	$1.0 \cdot 10^{-2}$	$2.6 \cdot 10^2$

(5.3)

In the first column we indicate the required accuracy. For each method, there are three subcolumns. First one indicates the number of iterations. Second one shows the upper estimate for the achieved accuracy.<sup>2</sup> The third column shows the current level of “Lipschitz constant”, generated by the method. Note that per one iteration of FGM we need in average to call the oracle four times. PGM needs in average only two calls.

It is clear, that both methods behave much better than the worst-case complexity bounds. For FGM, decrease of required accuracy in two times results in a doubling of the number of iterations. This dependence is typical for the complexity bounds of the type

---

<sup>2</sup>Since in this problem the optimal value is known, we use it in the stopping criterion.

$O(\frac{1}{\epsilon})$ , and not to the theoretical bound  $O(\frac{1}{\epsilon^2})$ . For PGM, the average increase is higher (approximately, in three times). But it is still much better than the theoretical bound.

From table (5.3) we can see that FGM generates much better model of the objective function. Its accuracy is usually only twice bigger than the actual residual. The estimates of PGM are much weaker. One of the possible reasons for this difference consists in much more aggressive behavior of FGM in introducing new gradients in the model.

Finally, the third subcolumn shows that the actual level of the Lipschitz constants are much lower than the theoretical prediction.

Let us look now how efficient FGM is in solving the same problem by the Euclidean setup. These results are presented in the left part of table (5.4).

$Eps$	FGM <sub>Euclid</sub>			WDA <sub>Entropy</sub>		
$2^{-5}$	886	$1.7 \cdot 10^{-2}$	$1.0 \cdot 10^6$	1569	$4.4 \cdot 10^{-2}$	1.0
$2^{-6}$	3249	$9.0 \cdot 10^{-2}$	$8.4 \cdot 10^6$	6086	$2.2 \cdot 10^{-2}$	1.0
$2^{-7}$	11803	$4.8 \cdot 10^{-2}$	$6.7 \cdot 10^7$	20655	$1.1 \cdot 10^{-2}$	1.0
$2^{-8}$	45417	$2.5 \cdot 10^{-2}$	$5.4 \cdot 10^8$	78832	$5.5 \cdot 10^{-3}$	1.0
$2^{-9}$	178866	$1.3 \cdot 10^{-2}$	$4.3 \cdot 10^9$	283352	$2.7 \cdot 10^{-3}$	1.0
$2^{-10}$	out of time			out of time		

(5.4)

They confirm that the right choice of prox-function is crucial for the efficient solution of optimization problems. Behavior of FGM with Euclidean setup just corresponds to the worst-case theoretical bound  $O(\frac{1}{\epsilon^2})$  for Lipschitz-continuous functions (increase of the number of iterations in four times after dividing accuracy by two).

In the right part of this table we present the results of the standard black-box subgradient scheme as applied to the same problem. This is Weighted Dual Averaging (WDA) [11] with Entropy Setup. For choosing its parameters correctly, we need to know only an estimate for the diameter of the feasible set. Each iteration of this method needs one call of oracle. For our problem, WDA works in an exact correspondence to its worst-case complexity bound  $O(\frac{1}{\epsilon^2})$ . The second column of this part demonstrates that the lower bound generated by this scheme is almost exact.

**2. Continuous Steiner problem.** In this problem we are given by centers  $a_i \in R^n$ ,  $i = 1, \dots, m$ . It is necessary to find the optimal location of the service center  $x$ , which minimizes the total distance to all other centers. Thus, our problem is as follows:

$$\min_{x \in Q} f(x) \stackrel{\text{def}}{=} \sum_{i=1}^m \|x - a_i\|. \quad (5.5)$$

where  $Q \subseteq R^n$  is a closed convex set. All norms in this problem are Euclidean.

Clearly, the level of smoothness of problem (5.5) is much higher than that of (5.2). So, we can expect that it is easier for the universal schemes. Let us look at the results of the experiments for random problem with  $n = 256$ ,  $m = 512$ , and  $Q = R_+^n$ . We choose  $m > n$  in order to increase the density of nonsmooth points. The centers were generated randomly in the box  $0 \leq x^{(i)} \leq \frac{1}{n^{1/2}}$ ,  $i = 1, \dots, n$  (which has Euclidean diameter one). All methods have origin as a starting point. The initial value of the objective is  $f_0 = 295.226$ . The optimal solution found by the schemes is  $f^* = 147.336$ . The table below has the same structure as (5.3).

$Eps$	FGM <sub>Euclid</sub>			PGM <sub>Euclid</sub>		
$2^{-5}$	205	$3.1 \cdot 10^{-2}$	$2.6 \cdot 10^2$	9925	$3.1 \cdot 10^{-2}$	$2.6 \cdot 10^2$
$2^{-6}$	307	$1.5 \cdot 10^{-2}$	$5.1 \cdot 10^2$	19895	$1.5 \cdot 10^{-2}$	$5.1 \cdot 10^2$
$2^{-7}$	277	$6.8 \cdot 10^{-3}$	$2.6 \cdot 10^2$	39803	$7.8 \cdot 10^{-3}$	$2.6 \cdot 10^2$
$2^{-8}$	611	$3.9 \cdot 10^{-3}$	$5.1 \cdot 10^2$	77138	$3.9 \cdot 10^{-3}$	$5.1 \cdot 10^2$
$2^{-9}$	827	$1.9 \cdot 10^{-3}$	$5.1 \cdot 10^2$	155038	$2.0 \cdot 10^{-3}$	$2.6 \cdot 10^2$
$2^{-10}$	1226	$9.8 \cdot 10^{-4}$	$2.6 \cdot 10^2$	out of time		
$2^{-11}$	1655	$4.8 \cdot 10^{-4}$	$2.6 \cdot 10^2$			
$2^{-12}$	2385	$2.4 \cdot 10^{-4}$	$5.1 \cdot 10^2$			
$2^{-13}$	3388	$1.2 \cdot 10^{-4}$	$5.1 \cdot 10^2$			

(5.6)

Note that the rate of convergence of FGM (4.1) is unexpectedly high. Increase of the accuracy in four times results in doubling the number of iterations. From the complexity point of view, this corresponds to the level  $O(\frac{1}{\epsilon^{1/2}})$ , which is typical for Fast Gradient Methods of *smooth* minimization. The predicted accuracy by FGM is still very good, and the level of Lipschitz constants is unexpectedly small. The results of PGM (2.16) are not so impressive. It doubles the number of iterations after dividing accuracy by two, which corresponds to  $O(\frac{1}{\epsilon})$  level of complexity. It seems that a weak point of this method is the quality of termination criterion.

**3. Universal methods and smoothing technique.** Let us compare the practical performance of method (4.1) as applied to the primal version of problem (5.1)

$$\min_{x \in \Delta_n} \psi_p(x), \quad (5.7)$$

with its performance as applied to the smoothed version of this function

$$\tilde{\psi}_p(x) = \mu \ln \left( \sum_{j=1}^m e^{\langle x, Ae_j \rangle / \mu} \right).$$

The value of smoothing parameter  $\mu > 0$  for this function is chosen in accordance to the theoretical recommendation (4.8) in [9]. For our experiments, we choose  $n = m = 512$ .

$Eps$	FGM - smoothing			FGM <sub>Entropy</sub>		
$2^{-5}$	47	$3.0 \cdot 10^{-2}$	$4.0 \cdot 10^0$	555	$3.1 \cdot 10^{-2}$	$1.0 \cdot 10^3$
$2^{-6}$	103	$1.5 \cdot 10^{-2}$	$8.0 \cdot 10^0$	1956	$1.5 \cdot 10^{-2}$	$1.6 \cdot 10^4$
$2^{-7}$	226	$7.6 \cdot 10^{-3}$	$1.6 \cdot 10^1$	8048	$7.8 \cdot 10^{-3}$	$2.6 \cdot 10^5$
$2^{-8}$	464	$3.9 \cdot 10^{-3}$	$3.2 \cdot 10^1$	34355	$3.9 \cdot 10^{-3}$	$1.0 \cdot 10^6$
$2^{-9}$	953	$1.9 \cdot 10^{-3}$	$1.3 \cdot 10^2$	135419	$2.0 \cdot 10^{-3}$	$8.4 \cdot 10^6$
$2^{-10}$	1881	$9.7 \cdot 10^{-4}$	$1.3 \cdot 10^2$	out of time		
$2^{-11}$	3653	$4.9 \cdot 10^{-4}$	$2.6 \cdot 10^2$			
$2^{-12}$	7077	$2.4 \cdot 10^{-4}$	$2.0 \cdot 10^3$			
$2^{-13}$	13771	$1.2 \cdot 10^{-4}$	$1.0 \cdot 10^3$			

(5.8)

These results confirm that smoothing is still a very powerful technique, which computational efficiency is often much higher than that of the Black Box Methods.

## References

- [1] A. Beck, M. Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. Imaging Sciences*, **2**(1), 183-202 (2009)
- [2] O.Devolder, F.Glineur, Yu.Nesterov. First-order methods of smooth convex optimization with inexact oracle. Accepted by *Mathematical Programming*.
- [3] K.-H.Elster(eds). *Modern Mathematical Methods in Optimization*. Academie Verlag, Berlin, 1993.
- [4] G. Lan. Level methods uniformly optimal for composite and structured nonsmooth convex optimization. April 2011, Submitted to *Mathematical Programming*.
- [5] C. Lemarechal, A. Nemirovskii and Yu. Nesterov. New variants of bundle methods. *Mathematical Programming*, 69 (1995), 111-147.
- [6] A. Nemirovskii, Yu. Nesterov. Optimal methods for smooth convex optimization. *Zh. Vychisl. Mat. i Mat. Fiz.* **25**(3), 356-369 (1985), in Russian.
- [7] A. Nemirovsky, D. Yudin. *Problem complexity and method efficiency in optimization* John Wiley & Sons, New York (1983).
- [8] Yu. Nesterov. *Introductory Lectures on Convex Optimization*. Kluwer, Boston, 2004.
- [9] Yu. Nesterov. "Smooth minimization of non-smooth functions", *Mathematical Programming (A)*, **103** (1), 127-152 (2005).
- [10] Yu.Nesterov. Gradient methods for minimizing composite functions. CORE DP 2007/76. Published in *Mathematical Programming*, December (2012)
- [11] Yu. Nesterov. "Primal-dual subgradient methods for convex problems". *Mathematical Programming*, **120**(1), 261-283 (August 2009)

## Recent titles

### CORE Discussion Papers

- 2012/49 Thierry BRECHET, Carmen CAMACHO and Vladimir M. VELIOV. Adaptive model-predictive climate policies in a multi-country setting.
- 2012/50 Vladyslav NORA and Hiroshi UNO. Saddle functions and robust sets of equilibria.
- 2012/51 Thomas BAUDIN, David DE LA CROIX and Paula GOBBI. DINKs, DEWks & Co. Marriage, fertility and childlessness in the United States.
- 2012/52 David DE LA CROIX and Omar LICANDRO. The longevity of famous people from Hammurabi to Einstein.
- 2012/53 Marijn VERSCHELDE, Jean HINDRIKS, Glenn RAYP and Koen SCHOORS. School staff autonomy and educational performance: within school type evidence.
- 2012/54 Thierry BRECHET and Susana PERALTA. Markets for tradable emission permits with fiscal competition.
- 2012/55 Sudipto BHATTACHARYA, Claude D'ASPREMONT, Sergei GURIEV, Debapriya SEN and Yair TAUMAN. Cooperation in R&D: patenting, licensing and contracting.
- 2012/56 Guillaume WUNSCH, Michel MOUCHART and Federica RUSSO. Functions and mechanisms in structural-modelling explanations.
- 2012/57 Simone MORICONI, Pierre M. PICARD and Skerdilajda ZANAJ. Commodity taxation and regulatory competition.
- 2012/58 Yurii NESTEROV and Arkadi NEMIROVSKI. Finding the stationary states of Markov chains by iterative methods.
- 2012/59 Tanguy ISAAC and Paolo PIACQUADIO. Equity and efficiency in an overlapping generation model.
- 2012/60 Luc BAUWENS, Giuseppe STORTI and Francesco VIOLANTE. Dynamic conditional correlation models for realized covariance matrices.
- 2012/61 Mikhail ESKAKOV and Alexey ISKAKOV. Equilibrium in secure strategies.
- 2012/62 Francis BLOCH and Axel GAUTIER. Strategic bypass deterrence.
- 2012/63 Olivier DURAND-LASSERVE, Axel PIERRU and Yves SMEERS. Sensitivity of policy simulation to benchmark scenarios in CGE models: illustration with carbon leakage.
- 2013/1 Pierre PESTIEAU and Maria RACIONERO. Harsh occupations, health status and social security.
- 2013/2 Thierry BRECHET and Henry TULKENS. Climate policies: a burden or a gain?
- 2013/3 Per J. AGRELL, Mehdi FARSI, Massimo FILIPPINI and Martin KOLLER. Unobserved heterogeneous effects in the cost efficiency analysis of electricity distribution systems.
- 2013/4 Adel HATAMI-MARBINI, Per J. AGRELL and Nazila AGHAYI. Imprecise data envelopment analysis for the two-stage process.
- 2013/5 Farhad HOSSEINZADEH LOTFI, Adel HATAMI-MARBINI, Per J. AGRELL, Kobra GHOLAMI and Zahra GHELEJ BEIGI. Centralized resource reduction and target setting under DEA control.
- 2013/6 Per J. AGRELL and Peter BOGETOFT. A three-stage supply chain investment model under asymmetric information.
- 2013/7 Per J. AGRELL and Pooria NIKNAZAR. Robustness, outliers and Mavericks in network regulation.
- 2013/8 Per J. AGRELL and Peter BOGETOFT. Benchmarking and regulation.
- 2013/9 Jacques H. DREZE. When Borch's Theorem does not apply: some key implications of market incompleteness, with policy relevance today.
- 2013/10 Jacques H. DREZE. Existence and multiplicity of temporary equilibria under nominal price rigidities.
- 2013/11 Jean HINDRIKS, Susana PERALTA and Shlomo WEBER. Local taxation of global corporation: a simple solution.
- 2013/12 Pierre DEHEZ and Sophie POUKENS. The Shapley value as a guide to FRAND licensing agreements.

## Recent titles

### CORE Discussion Papers - continued

- 2013/13 Jacques H. DREZE and Alain DURRE. Fiscal integration and growth stimulation in Europe.  
2013/14 Luc BAUWENS and Edoardo OTRANTO. Modeling the dependence of conditional correlations on volatility.  
2013/15 Jens L. HOUGAARD, Juan D. MORENO-TERNERO and Lars P. OSTERDAL. Assigning agents to a line.  
2013/16 Olivier DEVOLDER, François GLINEUR and Yu. NESTEROV. First-order methods with inexact oracle: the strongly convex case.  
2013/17 Olivier DEVOLDER, François GLINEUR and Yu. NESTEROV. Intermediate gradient methods for smooth convex problems with inexact oracle.  
2013/18 Diane PIERRET. The systemic risk of energy markets.  
2013/19 Pascal MOSSAY and Pierre M. PICARD. Spatial segregation and urban structure.  
2013/20 Philippe DE DONDER and Marie-Louise LEROUX. Behavioral biases and long term care insurance: a political economy approach.  
2013/21 Dominik DORSCH, Hubertus Th. JONGEN, Jan.-J. RÜCKMANN and Vladimir SHIKHMAN. On implicit functions in nonsmooth analysis.  
2013/22 Christian M. HAFNER and Oliver LINTON. An almost closed form estimator for the EGARCH model.  
2013/23 Johanna M. GOERTZ and François MANIQUET. Large elections with multiple alternatives: a Condorcet Jury Theorem and inefficient equilibria.  
2013/24 Axel GAUTIER and Jean-Christophe PODOU. Reforming the postal universal service.  
2013/25 Fabian Y.R.P. BOCART and Christian M. HAFNER. Fair re-valuation of wine as an investment.  
2013/26 Yu. NESTEROV. Universal gradient methods for convex optimization problems.

### Books

- G. DURANTON, Ph. MARTIN, Th. MAYER and F. MAYNERIS (2010), *The economics of clusters – Lessons from the French experience*. Oxford University Press.  
J. HINDRIKS and I. VAN DE CLOOT (2011), *Notre pension en héritage*. Itinera Institute.  
M. FLEURBAEY and F. MANIQUET (2011), *A theory of fairness and social welfare*. Cambridge University Press.  
V. GINSBURGH and S. WEBER (2011), *How many languages make sense? The economics of linguistic diversity*. Princeton University Press.  
I. THOMAS, D. VANNESTE and X. QUERRIAU (2011), *Atlas de Belgique – Tome 4 Habitat*. Academia Press.  
W. GAERTNER and E. SCHOKKAERT (2012), *Empirical social choice*. Cambridge University Press.  
L. BAUWENS, Ch. HAFNER and S. LAURENT (2012), *Handbook of volatility models and their applications*. Wiley.  
J-C. PRAGER and J. THISSE (2012), *Economic geography and the unequal development of regions*. Routledge.  
M. FLEURBAEY and F. MANIQUET (2012), *Equality of opportunity: the economics of responsibility*. World Scientific.  
J. HINDRIKS (2012), *Gestion publique*. De Boeck.

### CORE Lecture Series

- R. AMIR (2002), Supermodularity and complementarity in economics.  
R. WEISMANTEL (2006), Lectures on mixed nonlinear programming.  
A. SHAPIRO (2010), Stochastic programming: modeling and theory.